

学校编码: 10384

分类号_____密级_____

学 号: 200226005

UDC_____

厦 门 大 学
硕 士 学 位 论 文

不同群落类型长苞铁杉林根际
微生物研究

**Studies on the rhizospheric microorganisms of *Tsuga*
Longibracteata in different communities**

向 伟

指导教师姓名: 李 振 基 副教授

专 业 名 称: 植 物 学

论文提交日期: 2005 年 7 月 28 日

论文答辩日期: 2005 年 8 月 28 日

学位授予日期: 2005 年 月 日

答辩委员会主席: 郑文教 教 授

评 阅 人: 陈孝麟 研究员

庄铁城 教 授

2005 年 8 月

厦门大学学位论文原创性声明

兹提交的学位论文，是本人在导师指导下独立完成的研究成果。本人在论文写作中参考的其他个人或集体的研究成果，均在文中以明确方式标明。本人依法享有和承担由此论文而产生的权利和责任。

声明人（签名）：

年 月 日

厦门大学博研

目 录

摘 要.....	1
Abstract.....	1
第一章 前 言.....	1
1.1 长苞铁杉国内外研究概况.....	1
1.2 根际土壤微生物国内外研究概况.....	2
1.3 植物与根际土壤微生物相互作用研究进展.....	4
1.3.1 根际土壤微生物对植物生长的影响.....	5
1.3.2 植物根系对根际土壤微生物的影响.....	8
1.4 本实验的研究目的及意义.....	11
1.4.1 研究目的.....	11
1.4.2 研究意义.....	12
第二章 样地概况.....	13
2.1 自然条件概况.....	13
2.1.1 地质.....	13
2.1.2 地貌.....	13
2.1.3 气候条件.....	13
2.1.4 土壤条件.....	14
2.2 植物与植被资源.....	14
2.2.1 植物种类.....	14
2.2.2 植被概况.....	14
第三章 研究方法.....	16
3.1 土壤异养微生物(细菌、放线菌、丝状真菌)数量测定.....	16
3.2 林地中优势根际土壤异养微生物鉴定.....	16

3.2.1 细菌	16
3.2.2 放线菌	18
3.2.3 丝状真菌	18
3.3 不同群落根际、非根际土壤理化分析.....	19
第四章 结果与分析	20
4.1 微生物数量及根际效应.....	20
4.1.1 不同群落类型根际、非根际土壤异养微生物数量和组成	20
4.1.2 不同群落类型根际效应	21
4.2 菌种鉴定.....	24
4.2.1 细菌	24
4.2.2 放线菌	24
4.2.3 真菌	25
4.3 土壤理化性质.....	27
第五章 讨 论	29
5.1 微生物组成及数量.....	29
5.2 菌种鉴定	30
5.3 土壤理化性质.....	31
5.4 混交效应.....	33
第六章 结语	34
参考文献:	35
附录 1 天宝岩自然保护区地图.....	44
附录 2 样地微生物主要属种名录.....	45
附录 3 已发表论文目录.....	47
致 谢	48

CATALOG

Abstract in Chinese	I
Abstract in English	I
Chapter 1. Prolegomenon	1
1.1 Survey of <i>Tsuga longibracteata</i> Research At Home And abroad.....	1
1.2 Survey of rhizospheric microorganisms Research At Home And abroad.....	2
1.3 Advances of Interactions Between Plant and rhizospheric microorganisms.....	4
1.3.1 Rhizospheric Microorganisms's Affections to the Plant Growth.....	5
1.3.2 Plant Root's Affections to The Rhizospheric Microorganisms.....	8
1.4 The Objective and Significance of This Reserch.....	11
1.4.1 Objective.....	11
1.4.2 Significance.....	12
Chapter 2. Sample Field Instruction	13
2.1 Natural Conditions Instruction.....	13
2.1.1 Geology.....	13
2.1.2 Physiognomy.....	13
2.1.3 Climate conditions.....	13
2.1.4 Soil conditions.....	14

2. 2 Plangt and Vegetation resource	14
2. 2. 1 Plant specieses	14
2. 2. 2 Vegetation Instruction	14
Chapter 3. Methods	16
3. 1 Number of Soil microorganisms	16
3. 2 Identifiction of The Dominant Rhizospheric Microorganisms in Stands	16
3. 2. 1 Bacteria	16
3. 2. 2 Actinomycetes	18
3. 2. 3 Fungi	18
3. 3 The Analysis of Rhizospheric and Non-Rhizospheric Soil in Different Stands	19
Chapter 4. Result and Analysis	20
4. 1 number of Rhizospheric microorganisms and rhizospheric Effect	20
4. 1. 1 The Count and Compositions of Rhizospheric and Non- Rhizospheric Microorganisms in Different Stands	20
4. 1. 2 The Rhizospheric Effect in Different Stands	22
4. 2 Microorganisms Identification	24
4. 2. 1 Bacteria	24
4. 2. 2 Actinomycetes	24
4. 2. 3 Fungi	25
4. 3 The Physical and Chemical Characteristics of Soil	27

Chapter 5. Discussions	29
5. 1 Number and composition of microorganisms	29
5. 2 Microorganisms Identification	30
5. 3 The Physical and Chemical Characteristics of Soil	31
5. 4 effect of Mixed forest	33
Chapter 6. Epilogue	34
References	35
Appendix 1: Map of Tian-BaoYan natural reserve	44
Appendix 2: List of Microorganisms in Sample Field	45
Appendix 3: Publications during graduate study	47

厦门大学博研

摘 要

长苞铁杉是我国特有的珍稀古老植物，福建省省级保护植物，它材质优良，是南岭山脉乃至广大中亚热带中海拔山地极有潜力的珍贵用材树种和生态恢复树种。本实验以长苞铁杉(*Tsuga longibracteata*)群落(TLC)、长苞铁杉+猴头杜鹃(*Rhododendron simiarum*)群落(TRC)，长苞铁杉+毛竹(*Phyllostachys pubescens*)群落(TPC)根际土壤微生物为研究对象，调查不同群落类型单位重量根际土壤和非根际土壤中的异养微生物总数；三大类微生物(细菌、放线菌、真菌)各自的数量及其在总数量中所占的比例；不同群落的根际效应；根际土壤微生物种类和根际与非根际土壤理化性质。结果如下：

1. 三个群落类型中，微生物数量及组成各不相同(单位： $\text{cfu/g}\cdot\text{dw}\times 10^4$)：在长苞铁杉群落中：根际细菌(20.00)>非根际细菌(12.00)>根际真菌(9.37)>非根际真菌(6.74)>根际放线菌(2.69)>非根际放线菌(1.83)；在长苞铁杉+猴头杜鹃群落中：根际细菌(34.90)>非根际细菌(18.00)>根际真菌(6.54)>非根际真菌(3.78)>根际放线菌(2.96)>非根际放线菌(1.76)；在长苞铁杉+毛竹群落中：非根际细菌(88.00)>根际细菌(68.60)>非根际放线菌(13.40)>根际放线菌(12.60)>根际真菌(4.64)>非根际真菌(3.46)；
2. 通过比较根际与非根际土壤微生物得出不同长苞铁杉群落类型的根际效应，三大群落类型的总微生物的根际效应大小为：长苞铁杉+猴头杜鹃群落(1.87)>长苞铁杉群落(1.56)>长苞铁杉+毛竹群落(0.82)；
3. 从不同长苞铁杉群落的根际土壤中共分离到 8 种芽孢杆菌：蜡状芽孢杆菌(*Bacillus cereus*)、凝结芽孢杆菌(*Bacillus coagulans*)、巨大芽孢杆菌(*Bacillus megaterium*)、环状芽孢杆菌(*Bacillus circulans*)、球形芽孢杆菌(*Bacillus sphaericus*)、短小芽孢杆菌 (*Bacillus pumilus*)、短芽孢杆菌(*Bacillus brevis*)、枯草芽孢杆菌(*Bacillus subtilis*)；链霉菌属 7 个不同类群：灰褐类群(*Griseofuscus*)、烬灰类群(*Cinereus*)、青色类群(*Glaucus*)、白孢类群

(*Albosporus*)、金色类群(*Aureus*)、粉红孢类群(*Roseosporus*)、吸水类群(*Hygroscopicus*); 真菌 5 个属: 青霉属(*Penicillium*)、木霉属(*Trichoderma*)、毛霉属 (*Mucor*)、被孢霉属(*Mortierella*)、伞状霉属(*Umbelopsis*);

4. 各群落类型的 pH 值: 长群落非根际土壤(4.17)>长苞铁杉+猴头杜鹃群落非根际土壤(4.13)>长苞铁杉+毛竹群落非根际土壤(4.03)>长群落根际土壤(4.06)>长苞铁杉+毛竹群落根际土壤(3.94)>长苞铁杉+猴头杜鹃群落根际土壤(3.89);
5. 各群落类型的有机质(%): 长苞铁杉+毛竹群落非根际土壤(3.979)>长苞铁杉+猴头杜鹃群落根际土壤(3.855)>长苞铁杉+毛竹群落根际土壤(3.655)>长苞铁杉群落根际土壤(3.648)>长苞铁杉+猴头杜鹃群落非根际土壤(3.465)>长苞铁杉群落非根际土壤(3.394);
6. 各群落类型的全 N 含量(%): 长苞铁杉+毛竹群落非根际土壤(0.163)>长苞铁杉+毛竹群落根际土壤(0.149)>长苞铁杉+猴头杜鹃群落根际土壤(0.130)>长群落根际土壤(0.110)>长群落非根际土壤(0.069)>长苞铁杉+猴头杜鹃群落非根际土壤(0.062);
7. 各群落类型的全 P 含量(%): 长苞铁杉+毛竹群落非根际土壤(0.020)>长苞铁杉+猴头杜鹃群落根际土壤(0.017)>长苞铁杉+毛竹群落根际土壤(0.016)>长群落根际土壤(0.015)>长苞铁杉+猴头杜鹃群落非根际土壤(0.014)>长群落非根际土壤(0.013)。

关键词: 长苞铁杉; 根际土壤微生物; 根际效应。

Abstract

Tsuga longibracteata is a kind of rare and immemorial plant listed as the protected tree in Fujian Province and is endemic to China. *Tsuga longibracteata* has good properties and it is also one important kind of potential timber tree and for restoration in the mountains of southern China. The rhizospheric microorganisms in the stand of *Tsuga longibracteata* community (TLC), the stand of *Tsuga longibracteata* + *Rhododendron simiarum* community (TRC), and the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* community (TPC) in Tianbaoyan National Nature Reserve of Fujian were studied from 2004 to 2005. The result showed that:

1. The counts and composition of microorganisms in three stand are different (the unit is $\text{cfu/g} \cdot \text{dw} \times 10^4$): In the stand of *Tsuga longibracteata* association: rhizospheric bacterium (20.00) > non-rhizospheric bacterium (12.00) > rhizospheric fungi (9.37) > non-rhizospheric fungi (6.74) > rhizospheric actinomyces (2.69) > non-rhizospheric actinomyces (1.83); In the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association: rhizospheric bacterium (34.90) > non-rhizospheric bacterium (18.00) > rhizospheric fungi (6.54) > non-rhizospheric fungi (3.78) > rhizospheric actinomyces (2.96) > non-rhizospheric actionmyces (1.76); In the stand of *Tsuga longibracteata* and *Phyllostachys pubescens* association: non-rhizospheric bacterium (88.00) > rhizospheric bacterium (68.60) > non-rhizospheric actinomyces (13.40) > rhizospheric actinomyces (12.60) > rhizospheric fungi (4.64) > non-rhizospheric fungi (3.46).
2. The rhizospheric effect is: the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association (1.87) > the stand of *Tsuga longibracteata* association

- (1.56) > the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (0.82).
3. Eight *bacilluses* were isolated from the rhizospheric soil in the stand of *Tsuga longibracteata* association, the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association and the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association: *Bacillus cereus*, *Bacillus coagulans*, *Bacillus megaterium*, *Bacillus circulans*, *Bacillus sphaericus*, *Bacillus pumilus*, *Bacillus brevis*, *Bacillus subtilis*; seven *streptomyces* of actinomycetes: *Griseofuscus*, *Cinereus*, *Glaucus*, *Albosporus*, *Aureus*, *Roseosporus*, *Hygroscopicus*; Five major genera of the fungi: *Penicillium*, *Trichoderma*, *Mucor*, *Mortierella*, *Aspergillus*.
4. The pH in three associations were: Non-rhizospheric soil in the stand of *Tsuga longibracteata* association (4.17)> Non-rhizospheric soil in the stand of *Tsuga longibracteata* and *Rhododendron simiarum* association (4.13)> Non-rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (4.03)> Rhizospheric soil in the stand of *Tsuga longibracteata* association (4.06)> Rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association(3.94)> Rhizospheric soil in the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association(3.89).
5. The organic matter(%) in three associations were: Non-rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (3.979) > Rhizospheric soil in the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association (3.855) > Rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (3.655) > Rhizospheric soil in the stand of *Tsuga longibracteata* association (3.648) >

Non-Rhizospheric soil in the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association (3.465) > Non-rhizospheric soil in the stand of *Tsuga longibracteata* association(3.394).

6. The contents of N(%) in three associations were: Non-rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (0.163) > Rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (0.149) > Rhizospheric soil in the stand of *Tsuga longibracteata* and *Rhododendron simiarum* association (0.130) > Rhizospheric soil in the stand of *Tsuga longibracteata* association (0.112) > Non-rhizospheric soil in the stand of *Tsuga longibracteata* association (0.069) > Non-Rhizospheric soil in the stand of *Tsuga longibracteata* and *Rhododendron simiarum* association (0.062).
7. The contents of P(%) in three associations were: Non-rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (0.020) > Rhizospheric soil in the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association (0.017) > Rhizospheric soil in the stand of *Tsuga longibracteata* + *Phyllostachys pubescens* association (0.016) > Rhizospheric soil in the stand of *Tsuga longibracteata* association (0.015) > Non-Rhizospheric soil in the stand of *Tsuga longibracteata* + *Rhododendron simiarum* association (0.014) > Non-rhizospheric soil in the stand of *Tsuga longibracteata* association (0.013).

Key words: *Tsuga longibracteata*; rhizospheric microorganisms; rhizospheric effect.

厦门大学博研

第一章 前 言

1.1 长苞铁杉国内外研究概况

长苞铁杉是我国特有树种，20 世纪 90 年代以来，国内学者开始重视对它的研究。

在群落生态学方面，邹惠渝和周晓白(1994)通过对福建几处长苞铁杉林的天然更新研究，初步揭示了火生态促进天然更新的机理^[1]。王新功等(2003)对不同区域的长苞铁杉群落植物组成及其区系特征进行了比较研究^[2]。

在种群生态学方面，吴继林等(1999)首次将连续型 Weibull 分布用于长苞铁杉的种群空间分布格局的研究，效果明显，根据 Weibull 分布建立了长苞铁杉种群最适量抽样式，为长苞铁杉种群抽样技术和取样数量确定提供了理论依据^[3]。吴承祯等(2001)对长苞铁杉的空间分布格局进行了研究，得出长苞铁杉的空间分布呈随机分布^[4]。吴承祯等(2001)通过用重要值百分数求取种间竞争系数，采用 Lotka-Volterra 竞争方程研究长苞铁杉的种间竞争关系，说明了在长苞铁杉种群支配整个群落^[5]。

在森林水文学方面，钟祥顺(1999)比较了长苞铁杉天然林、常绿阔叶林和杉木人工林的林冠层、林下植被层和凋落物层的持水量、土壤蓄水能力和渗透性能，表明长苞铁杉天然林具有较好的水源涵养功能^[6]。

在植物资源学方面，李维林等(2001)对长苞铁杉枝叶的挥发油成分进行了相关研究^[7]。

综上所述，可以看出到目前为止国内对长苞铁杉的研究主要集中在长苞铁杉群落空间分布格局及其种群分布等方面，而对于长苞铁杉群落根际土壤微生物方面还鲜有研究。

长苞铁杉作为我国特有树种，只零星的见于我国福建、江西、广西、广

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

廈門大學博碩士論文